



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01K 67/027, C12N 5/06, 5/10	A1	(11) International Publication Number: WO 99/53751 (43) International Publication Date: 28 October 1999 (28.10.99)
(21) International Application Number: PCT/EP99/02624 (22) International Filing Date: 19 April 1999 (19.04.99) (30) Priority Data: 9808325.6 20 April 1998 (20.04.98) GB (71) Applicant (for all designated States except US): CONSORZIO INCREMENTO ZOOTECHNICO S.R.L. [IT/IT]; Via Porcellasco, 7-f, I-26100 Cremona (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): GALLI, Cesare [IT/IT]; Via Persico, 191/G, I-26100 Cremona (IT). LAZZARI, Giovanna [IT/IT]; Via Persico, 191/G, I-26100 Cremona (IT). (74) Agent: WAKERLEY, Helen, Rachael; Reddie & Grose, 16 Theobalds Road, London WC1X 8PL (GB).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: SOURCE OF NUCLEI FOR NUCLEAR TRANSFER (57) Abstract <p>The reconstruction of a mammalian embryo uses lymphocytes as the source of donor nuclei. The recipient may be an enucleated oocyte. The embryo so prepared may be brought to term, used in recloning techniques or used to prepare embryonic stem cell lines.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LJ	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

- 1 -

Source of nuclei for nuclear transfer

This invention relates to the generation of animals genetically identical to an existing or existed animal. Further, during the process of regeneration some characteristic(s) can be changed by recombinant DNA technology to produce a transgenic animal by the addition or deletion of selected genes.

Known procedures for nuclear transfer involve the transfer of a nucleus taken from a pre-implantation stage embryo into an enucleated mature oocyte. Following activation of the oocyte, in a process that mimics sperm entry and signalling, an embryo develops and eventually an individual that is genetically (as far as DNA is concerned) identical to the donor embryo. The limited number of cells present in a mammalian pre-implantation embryo, however, allows the regeneration of a limited number of embryos. Pre-implantation embryo nuclei donors do not allow the use of recombinant DNA technology because of the limited number of cells available. Most importantly, though, the genetic value of the embryo, and thus of the animal that will be born, can only be estimated. This is of low economic value. For these reasons the potential of nucleus transfer technology has not been developed with commercial exploitation in mind; its major use is for scientific purposes.

A partial solution to the limited number of nuclei has been the use of a so called 'serial nucleus transfer' where the embryos obtained from the starting embryos are further subjected once, or more than once, to the same procedure therefore increasing the number of embryos regenerated (Stice & al., 1991, Theriogenology 35, 273).

The major limitations to the use of nucleus transfer procedure outlined above would be overcome if a renewable and / or unlimited source of nuclei to be used in the process could be made available. For many years people have attempted to establish cell lines from pre-implantation embryos (embryonic stem cell lines) but failed except for the mouse. Such work is reviewed in Galli et al. 1994, Zygote 2: 385-389. This type of cell would represent the ideal source of nuclei,

CONFIRMATION COPY

- 2 -

however in the mouse they have never been used in nucleus transfer experiments.

Cultured inner cell mass cells or presumptive embryonic cell lines have been obtained and successfully used for nucleus transfer experiments to produce embryos: Moor, Sun & Galli, 1992, Animal Reprod. Sci. 28, 423-431; Stice & al. 1996, Biol. Reprod. 54, 100-110. Viable offspring have been produced in cattle and sheep: Sims & First, 1994, Proc. Natl. Acad. Sci. USA 91: 6143-6147; Campbell & al. 1996, Nature 380, 64-66; Wells & al. 1997, Biol. Reprod. 57, 385-393. More recently, viable offspring has also been obtained with the use of nuclei from cultured fetal cells: Wilmut & al. 1997, Nature 385, 810-813; see The New York Times 21 January 1998 and Nature 392, 113, 1998. One lamb has been produced from a sample taken from a primary culture containing mainly mammary epithelial cells of an adult sheep: Wilmut & al. 1997, Nature 385, 810-813.

The advantages of using a renewable source of cell or a cell line in nucleus transfer procedure are:

- cells can be easily collected and cultivated or possibly stored in liquid nitrogen;
- an unlimited number of embryos could be produced over a long period;
- cells can readily be modified in vitro using recombinant DNA technology.

There has been discussion about using nuclei of somatic cells collected from adult animals. This will have particular application for livestock species where the value of an animal is determined by his progeny if a sire or by her production records if, for example, a dam. To regenerate a unique animal for production or genetic characteristics (transgenic) it is imperative to use nuclei from an animal which is an adult or one which has at least been born alive. That is not the case for the work using fetal or embryonic cells as a source of nuclei.

- 3 -

Description of the invention

The present invention provides a method of reconstructing a mammalian embryo, the method comprising transferring a lymphocyte into a suitable recipient.

The lymphocyte can be transferred intact, optionally with a broken cell membrane, or the nucleus may be extracted and used for transfer. Preferably, the lymphocyte is transferred with the cell membrane broken.

Preferably, this invention finds application in the reconstruction of embryos of mammals using donor cells and recipients from the same species, preferably to reconstruct ungulate species embryos. The lymphocyte may be collected from an adult animal or an animal from a viable birth.

The invention further provides a method of reconstructing a mammalian embryo comprising reconstructing a first generation embryo by the steps of a method according to the first aspect of the invention and then transferring a cell from the said first generation embryo to a suitable recipient to form a second generation embryo.

The invention still further provides a method of preparing a mammal, the method comprising reconstructing a mammalian embryo using a method described above; allowing the embryo so produced to develop to term; and, optionally, breeding from the animal so formed.

The present invention further provides a method of preparing embryonic stem cell lines, comprising reconstructing an animal, preferably mammalian embryo using a method described above; and transferring the embryo to a culture system.

The present invention further provides a method of preparing embryonic stem cell lines, comprising reconstructing an animal, preferably mammalian embryo using a method described above; isolating the inner cell mass of the embryo from the embryo; and transferring the inner cell mass to a culture system.

- 4 -

The culture system allows the embryo cells to attach, outgrow and produce a cell line with embryonic characteristics. The term "embryo" used herein includes morulas (8-16 cells), morulas (16-32 cells) and blastocysts (64 cells and above). The embryo has a reasonable (about 50% or more) chance of development to an established pregnancy.

The present invention utilises lymphocytes and their derivatives or precursors. The donor cell, whilst usually a terminally differentiated hematopoietic cell, could be at a partially differentiated stage. These are mononuclear cells of hematopoietic lineage, present in bone marrow, lymphoid organs and in peripheral blood. They are also found in the umbilical cord of the new-born. The intact cells, cells with their membranes broken prior to transfer or the isolated nuclei are used as a source of nuclei in conventional nucleus transfer procedures. Lymphocytes can be collected from circulating blood, bone marrow, cord blood, lymphoid organs or natural secretions including milk and ejaculated semen. The sample can be enriched and purified by means of density gradient centrifugation or other means of separation, including immunomagnetic separation, fluorescence activated cell sorting, column filtration and similar techniques.

In the context of this invention, references to "mononuclear cells" for donor cells should be interpreted as references to lymphocytes, being lymphocytes at more than 95% of the mononuclear cells separated on a Hystopaque gradient. Lymphocytes have been characterised by immunocytochemistry and do not express cytokeratins as well as lamin A/C that are typical of differentiated cells: Galli & al. 1995, Proc. of the Italian Soc. of Vet. Sci. XLIX, 303-304; Rober, RA & al., 1990, J. Cell Sci. 95, 587-598. To this extent, the hematopoietic lineage shares some characteristics with embryonic cells that are also negative for cytokeratins and lamin A/C: Galli et al. 1994, Zygote 2: 385-389. This could explain in part the successful reprogramming of these nuclei into the cytoplasm of enucleated matured oocytes.

Freshly collected lymphocytes can be cultured in vitro and are karyotypically normal. This latter characteristic is a prerequisite for the normal development of any individual, but it is not guaranteed by other cell types that have to be cultured for a length of time and

- 5 -

where a degree of aneuploidy always occurs. Lymphocytes can also be cultured in vitro for a time sufficient to use recombinant DNA technology to alter their genetic constitution: Bordignon & al., 1995, Science 270, 470-475.

In principle this invention is applicable to all animals, but it will be useful in particular for livestock species such as cattle, buffaloes, sheep, goat, pigs, horses, rabbit and other species of economic relevance. It can also be used to preserve genetic material or to generate animals of endangered, exotic or rare species. In humans, it could find beneficial application in its use to generate embryonic stem cells from a patient as a source of compatible undifferentiated cells to be used in transplantation for the therapy of degenerative diseases.

After the reconstruction procedure whereby a lymphocyte or the nucleus of a lymphocyte is reprogrammed into the cytoplasm of an enucleated oocyte, there are several options for which this invention could be used. Lymphocytes can be easily cryobanked and therefore offer an economic way of storing germplasm of animals. When the embryo is reconstructed it can be used not for reproduction but to generate undifferentiated embryonic cell lines to be used in cell therapy of the individual that donated them thus overcoming the problem of rejection. If the embryos obtained are used for the generation of an animal this can be done directly by transferring the pre-implantation embryos to a final recipient that will carry the embryo to term, or the embryo can be subjected to serial nucleus transfer and therefore generate further embryos in a process that is more efficient and probably will increase the chances of reprogramming the cell nucleus because is exposed to the egg's cytoplasm more than once in a short period.

The steps involved in the cloning of an animal using this invention are summarised:

Step 1 - isolate the donor cell required from circulating blood or other tissue; enrichment for the fraction of cells that is more efficient in the procedure; optionally the cells can be genetically

- 6 -

modified during a period of in vitro culture using recombinant DNA technology.

At this stage the cells can be cultured, cryopreserved following one of the established protocols for later use or used immediately for nucleus transfer.

Step 2 - maturation of the oocytes harvested from donor females at slaughter or from live donors and removal of the egg's metaphase plate to prepare the so called 'recipient cytoplasm'.

Step 3 - transfer of the nucleus obtained in step 1 by direct microinjection of the cell or of the isolated nucleus directly in the cytoplasm of the enucleated oocyte or by other means such as cell fusion that can be achieved using intact donor cells with chemical, electrical or viral means. Microinjection is preferred and, preferably, the cell is transferred with the cell membrane broken. Established cell fusion methods include the use of fusion-promoting chemicals, such as polyethylene glycol; the use of a virus such as the Sendai virus; and electrical stimulation.

After introduction of the lymphocyte, the oocyte is activated to mimic sperm entry and start the developmental programme of the oocyte. The delay if microinjection is used to introduce the lymphocyte is typically 2-6 hours before activation. Cold shock as well as aging can activate the cytoplasm. Activation may also be by inducing calcium oscillations in the embryo by chemical (ionophore) or physical (electric current) means, following which the embryo is exposed to kinases and protein synthesis inhibitors that facilitate the exit from the metaphase arrest that is maintained upon new protein synthesis. Typical chemical activation would be by 6-dimethylamino purine or cycloheximide. This exposure would be subsequent to the ionophore and the exposure is typically for several hours (e.g., 4-6 hours).

Step 4 - develop the reconstructed embryo to a stage where it can be transferred to the uterus of the final recipient or subject to a serial cloning procedure by disaggregating the embryos obtained in single cells

- 7 -

and restarting from step 2. Various known systems of culturing embryos can be used successfully.

The steps involved in the preparation of a stem cell line using this invention are summarised:

Obtain a preimplantation stage (morula or blastocyst, preferably a blastocyst) embryo following steps 1-4 described in the previous example.

Step 5 - Remove the zona pellucida of the embryo. Optionally, the inner cell mass may be isolated from the embryo, for example by mechanical means or by immunosurgery. The intact embryo or the isolated inner cell mass is plated and cultured. Various known systems of culturing embryonic stem cells may be used. The culture takes place on a monolayer of fibroblasts and/or in defined media supplemented with the necessary growth factors (leukaemia inhibitor factor, stem cell factor and others), which are required to maintain the embryonic cell in an undifferentiated state.

Step 6 - Subculture using, for example, mechanical or enzyme dispersal of the embryonic cell outgrowths in new culture vessels to expand the number of cells until a stable cell line is obtained.

Step 7 - The cell line may be frozen for long term storage or the genetic constitution of the cells genetically modified using recombinant DNA technology.

Step 8 - Following genetic modification, the embryonic cell may be used in the cloning of a mammal by following steps 2 to 4.

Recloning procedures can also be carried out by developing the embryo of steps 1-4 to the fetus stage in vivo and sampling cells from the fetus for use in the preparation of further embryos.

EXAMPLE

This is an example of the use of the invention for the cloning of a cattle but similarly it can be applied to other species.

Step 1 - Cell isolation

A blood sample was taken from a cow of proven genetic value by venipuncture with heparinized vacutainer. The blood was diluted 1:1 with phosphate buffer saline (PBS) and 7 ml were layered on 3 ml of a density gradient (Hystopaque density 1083 g/cm³, Sigma) and centrifuged at 1500g for 15-30 minutes, the mononuclear cells stopping at the plasma Hystopaque interface. The 0.5 - 1 ml band of mononuclear cells (lymphocytes) was recovered, transferred into a new centrifuge tube, further diluted with PBS and centrifuged again to wash the cells. This step was repeated once and the cells were finally resuspended in an appropriate culture medium.

Lymphocytes were cryopreserved in medium supplemented with 10-20% serum and 10% DMSO (dimethyl sulfoxide) and packed for example in plastic straws (normally used to pack bovine semen), each containing convenient working aliquots of cells (0.5-2 million cells) required in each day the method of the invention was carried out.

Step 2 - Preparation of cytoplasts

Oocytes at the second metaphase were used. These oocytes were collected from ovaries of slaughtered animals or by ultrasound guided transvaginal recovery from live donors. After collection immature oocytes were subjected to a 15-20 hour maturation period until they reached the second metaphase, following protocols described by Galli & Lazzari, Anim. Reprod. Sci. 42, 371-379, 1996. Oocytes at the end of the maturation period were denuded from the surrounding follicle cells and treated with a fluorescent dye (Hoechst 33342) that stains the chromosomes in the metaphase plate. With the aid of a micromanipulator under an inverted microscope using a micropipette, the first polar body, with a small volume of cytoplasm surrounding it, was removed and checked under fluorescent light for the presence of the metaphase plate. After enucleation, the cytoplasts obtained in this way were returned to culture.

Step 3 - Embryo reconstruction

Lymphocytes prepared in step 1 and cytoplasts prepared in step 2 were transferred to a manipulation chamber under an inverted microscope and each cytoplast was injected with a small micropipette with one cell as described in Tesarik & Mendoza, Human Reproduction 11, 772-779; 1996. It was important to make sure that the cytoplast membrane was broken and the cell or its nucleus was effectively injected in the cytoplasm ready to undergo the reprogramming events necessary to support embryonic development. Failure to break the cytoplast membrane adequately could leave the lymphocyte deposited in a "pocket" of the oocyte membrane. After injection the cytoplasts were returned to culture for a period generally of 2-4 hours.

About 70-80% of the cytoplasts survived the injection procedure. At this stage the oocytes were activated by exposing sequentially the reconstructed embryos (cytoplasts) for 5-7 minutes to 5 μ M of Ionomycin (Sigma) and then to 2.5 mM 6-DMAP (Dimethyl amino purine, Sigma) for 4-5 hours: Susko-Parrish & al. 1994, Dev. Biol. 166, 729-739. This mimics sperm entry and will start the developmental programme of the oocyte.

Step 4 - Embryo development

Following activation, the reconstructed embryos were transferred to an in vitro culture system generally used to develop fertilised oocytes to blastocysts. Embryos were cultured in microdrops of SOF (synthetic oviductal fluid, Gardner & al. 1994; Biol. Reprod, 50, 390-400) in an atmosphere of 5% CO₂, 5%O₂ in nitrogen at 38.5 °C.

A proportion of the embryos (5%) developed to the blastocyst stage and could therefore be transferred to synchronised recipients or frozen for subsequent transfer.

With such embryos a pregnancy rate of over 50% (58%) was achieved, the most advanced stage obtained was a pregnancy aborted at 195 days of gestation (a normal bull calf of about 10 kg). The results are shown in Table 1. Most of the pregnancies resulted in abortions between 60-

- 10 -

120 days. Two pregnancies developed to 180 days from 31 transfers. The first is the 195 day mentioned above, the second an oversized bull calf of 19kg which had to be aborted.

Second Generation Cloning

In the first generation cloning only 5% of the reconstructed embryos developed to the blastocyst stage. By this, 16 cell stage by day 4 and compacted morula by day 6 are presumably those embryos in which the reprogramming of the introduced nucleus has occurred.

To increase the efficiency of the procedure the first generation products are subjected to a second generation cloning. This second generation cloning is more efficient because it uses blastomeres (16, 32, 64 cells stages, preferably 32 or over cells stage) and also gives the DNA a second chance for reprogramming because it is recycled back into the cytoplasm.

Embryos obtained in the first generation cloning were exposed to calcium and magnesium free HBSS (Hanks balanced salt solution) for 2-4 hours to separate the embryo into single isolated blastomeres.

Cytoplasts were prepared as described in step 2 and first activated (as described in step 3) before the blastomere nucleus was transferred. In this case, the intact blastomere was transferred to the perivitelline space of the cytoplasm and electrofused. The fusion rate was usually high (in excess of 80%). Reconstructed embryos at this stage were transferred to the culture system described above. The results are shown in Table 1. 19 such recloned embryos were transferred and 10 pregnancies were established, a pregnancy rate of over 50% (53%).

A successful birth has been achieved; a live and healthy calf originating from an embryo of the second generation cloning. The original lymphocyte was collected from a bull.

- 11 -

TABLE 1

Embryo development following lymphocyte injection or recloning.

	No. replicates	No. injected	No. survived %	No. cleaved %	No. developed %
Direct injection (development to blastocyst)	25	1923	1377 71.61	1059 76.91	71 5.16
Direct injection (development to morula 16-64 cells)	7	540	371 38.70	296 79.78	36 9.70
Recloned from morula (development to blastocyst)	7	462*	412 89.18	321 77.91	66 16.02

* "No. injected" is "No. fused" because recloning from blastomeres requires cell fusion not direct injection

Pregnancies	Totals	Cloning	Recloning
No. transfers	50	31	19
No. pregnancies	28	18	10
Pregnancy rate (%)	56.00	58.06	52.63
Developed to 90 days	10	5	5
Developed to 180 days	3	2	1
Developed to term	1	0	1

EFFECT OF THE INVENTION

The present invention provides a source of donor cells for nuclear transfer techniques which gives advantages over known donors. The use of lymphocytes makes for very easy sample collection, which can be from adult animals of known characteristics. The supply of donor cells is not limited. The donor cells can be readily modified in vitro using recombinant DNA technology.

Claims

1. A method of reconstructing a mammalian embryo, the method comprising transferring a lymphocyte into a suitable recipient.
2. The method according to claim 1 further comprising the step of isolating the nucleus of the lymphocyte before transfer of said nucleus into the recipient.
3. Method according to claim 1 or 2 in which the mammal is an ungulate species.
4. Method according to any preceding claim further comprising the step of genetically modifying the nucleus of the lymphocyte.
5. Method according to any preceding claim in which the recipient is an enucleate oocyte.
6. A method of reconstructing a mammalian embryo comprising reconstructing a first generation embryo by the steps of a method according to any of claims 1 to 5 and further comprising transferring a cell from the said first generation embryo to a suitable recipient to form a second generation embryo.
7. A method of reconstructing a mammalian embryo comprising reconstructing a first generation fetus by development of a first generation embryo reconstructed by a method of any of claims 1 to 5, preparing fetal fibroblast cultures therefrom and transferring cells from the said fetal fibroblast cultures to a suitable recipient to form a second generation embryo.
8. A method according to claim 7 further comprising the step of genetic modification of the cells of the fetal fibroblast cultures prior to second generation cloning.
9. A method of preparing a mammal, the method comprising:
reconstructing a mammalian embryo using a method according to any preceding claim;

- 14 -

allowing the embryo so produced to develop to term; and optionally breeding from the mammal so formed.

10. A method of preparing embryonic stem cell lines, comprising reconstructing a mammalian embryo using a method according to any of claims 1 to 8 and transferring the embryo to a culture system.

11. A method of preparing embryonic stem cell lines, comprising reconstructing a mammalian embryo using a method according to any of claims 1 to 8; isolating the inner cell mass of the embryo from the embryo and transferring the inner cell mass to a culture system.

12. A method according to claim 10 or 11 further comprising the step of genetic modification of the stem cells.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/02624

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A01K67/027 C12N5/06 C12N5/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 07841 A (UNIV MASSACHUSETTS) 26 February 1998 (1998-02-26)	1,2,5,6, 10-12
Y	* page 1, line 4-18, page 8, line 15-18, Table 1 *	3,4,7-9
Y	--- SCHNIEKE A ET AL: "Human Factor IX transgenic sheep produced by transfer of nuclei from transfected fetal fibroblasts" SCIENCE, vol. 278, 19 December 1997 (1997-12-19), pages 2130-2133, XP002067036 abstract	3,4,7-9
P,X	--- WO 98 30683 A (UNIV MASSACHUSETTS A PUBLIC IN) 16 July 1998 (1998-07-16) page 23, line 17 --- -/--	1-12

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

20 September 1999

Date of mailing of the international search report

05/10/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lonnoy, 0

INTERNATIONAL SEARCH REPORT

Intern: 1st Application No
PCT/EP 99/02624

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97 07668 A (CAMPBELL KEITH HENRY STOCKMAN ; ROSLIN INST EDINBURGH (GB); WILMUT) 6 March 1997 (1997-03-06) page 8, line 13-19 ---	
A	RITCHIE W A ET AL: "INTRACITOPLASMIC NUCLEAR INJECTION AS AN ALTERNATIVE TO CELL FUSION FOR THE PRODUCTION OF BOVINE EMBRYOS BY NUCLEAR TRANSFER" JOURNAL OF REPRODUCTION AND FERTILITY. SUPPLEMENT, vol. 5, 1 January 1995 (1995-01-01), page 60 XP000607293 ---	
A	DU PASQUIER L ET AL: "Transplantation of nuclei from lymphocytes of adult frogs into enucleated eggs: special focus on technical parameters" DIFFERENTIATION, vol. 8, no. 1, 1977, pages 9-19, XP002115398 abstract ---	
A	WO 97 07669 A (ROSLIN INST EDINBURGH ; CAMPBELL KEITH HENRY STOCKMAN (GB); WILMUT) 6 March 1997 (1997-03-06) ---	
P,A	KATO Y ET AL: "Eight calves cloned from somatic cells of a single adult" SCIENCE, vol. 282, no. 5396, 11 December 1998 (1998-12-11), pages 2095-2098, XP002115305 -----	

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 99/02624

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Claims 1, 2 and 4-12 (all partially) have not been searched in so far the embryo is a human embryo, as this subject matter falls within the exceptions to patentability of Article 53 (a) EPC.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/02624

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9807841 A	26-02-1998	AU 4044397 A	06-03-1998
		EP 0934403 A	11-08-1999
WO 9830683 A	16-07-1998	AU 6014598 A	03-08-1998
WO 9707668 A	06-03-1997	AU 6830996 A	19-03-1997
		CA 2229657 A	06-03-1997
		CN 1202085 A	16-12-1998
		CZ 9800604 A	15-07-1998
		EP 0847237 A	17-06-1998
		GB 2318792 A	06-05-1998
		HU 9802485 A	01-02-1999
		NO 980846 A	29-04-1998
		PL 325336 A	20-07-1998
WO 9707669 A	06-03-1997	AU 6831096 A	19-03-1997
		CA 2229568 A	06-03-1997
		CN 1202084 A	16-12-1998
		CZ 9800608 A	15-07-1998
		EP 0849990 A	01-07-1998
		EP 0930009 A	21-07-1999
		GB 2318578 A	29-04-1998
		GB 2331751 A	02-06-1999
		HU 9900234 A	28-05-1999
		NO 980845 A	29-04-1998
		PL 325331 A	20-07-1998